

Form field theory on Calabi-Yau threefold

Farhang Loran¹, Hesam Soltanpanahi²

*Department of Physics, Isfahan University of Technology,
Isfahan 84156-83111, Iran*

Abstract

We show how the natural Abelian duality of 2 and 3-form gravity theories on seven-dimensional manifold $CY_3 \times S^1$ leads to an S-duality between 2 and 3-form theories on simply connected CY_3 . The massless sector of the 2-form field theory on CY_3 corresponds to the A model string field theory. We discuss the complex structure independence of the 2 form theory for a general Kähler manifold and derive the holomorphic anomaly equations for simply connected CY_3 .

¹e-mail: loran@cc.iut.ac.ir

²e-mail: h_soltanpanahi@ph.iut.ac.ir

Contents

1	Introduction and conclusion	1
2	Abelian duality in 7 dimensions	4
2.1	Abelian duality	4
2.2	7-dimensional Abelian duality	5
2.3	Equation of motion and gauge fixing	8
3	Reduction to Calabi-Yau threefold	10
4	Canonical quantization and position/momentum duality	11
4.1	S-duality on CY_3	12
5	Theory of Kähler gravity	13
6	Quantum background independence	16
6.1	The structure of connections	17
6.2	Complex structure independence in form field theories	17
A	Abelian duality in 7 dimensions	19
B	Relation between the operators d and $*$ in m and $m + 1$ dimensions	19

1 Introduction and conclusion

Since the beginning of 90s, form theories of gravity have been studied by several authors. Interesting examples of them, related to the present work are the Chern-Simons theory exposition of 3-dimensional gravity [1], the description of the target space theory of A model topological string in terms of variations of the Kähler 2-form [2], and the description of the target space of B model topological string theory in terms of variations of the holomorphic 3-form [3]-[9].

In [10] Vafa conjectured that there is a mirror symmetry between topological A and B string theories. For topological string theories targeting a simply connected Calabi-Yau threefold, the observables of the A model are in one to one correspondence with the elements of $H^{1,1}(CY_3)$ while they are independent of the complex structure moduli. On the other hand, for the topological B model the observables are in one to one correspondence with the elements of $H^{2,1}(CY_3)$ but independent of the Kähler structure. The existence of such theories is supported by the Candelas observation in [11] that the moduli space of CY_3 factorizes into a direct product of complex structure moduli and Kähler moduli at least locally. For a simply connected Calabi-Yau threefold, $h^{1,1}$ and $h^{2,1}$ i.e. the dimensions of $H^{1,1}$ and $H^{2,1}$ respectively are the only nontrivial independent Hodge numbers. A Calabi-Yau threefold mirror to a given Calabi-Yau threefold is identified with Hodge numbers,

$$\tilde{h}^{1,1} = h^{2,1}, \quad \tilde{h}^{2,1} = h^{1,1}. \quad (1)$$

The conjectured mirror symmetry of [10] is a duality between the A model on a CY_3 and the B model on the corresponding mirror which has been confirmed for example for quintics and the corresponding mirrors. See e.g. [12] and references therein.

Roughly speaking the relation between the A model and B model topological string theories resembles the relation between type IIA and IIB superstring theories. The T-duality between type IIA and IIB superstring theories is explained in terms of an M-theory defined on the superstring target space trivially fibred over an extra Euclidean S^1 . Therefore it is natural to expect that there exists a topological M-theory in 7 dimensions which can be used to describe the duality of topological string field theories. Also, one expects that this 7-dimensional theory is defined on $CY_3 \times S^1$, where CY_3 is trivially fibred over S^1 . The trivially fibred $CY_3 \times S^1$ is one of the simplest examples of G_2 manifolds which are the natural classes of 7-dimensional manifolds preserving some supersymmetry and are purely geometric [13]. This is the setup of [14] where, a duality of A and B model string field theories on the same target space is studied. In principle, the mirror symmetry of the A and B string field theories can be given by a topological M-theory for which the CY_3 is non trivially fibred over $S^1/\mathbb{Z}_2$ which fixed points are the locations of a Calabi-Yau threefold and the corresponding mirror.

The topological A and B string field theories can be considered as form gravity theories in 6 or 7 dimensions. Recently, in [15, 16] it is conjectured that there is an S-duality between A and B models targeting the same CY_3 . They have shown that the perturbative amplitudes of the B model capture the non-perturbative amplitudes of the A model and vice versa. In [14] it is demonstrated that there is a 7-dimensional form theory on $CY_3 \times S^1$, called topological M-theory, which is related to both 2 and 3-form theories on the same CY_3 , and at least at the classical level these form theories are related to the topological A and B models respectively. Also, they have deduced a position/momentum duality, called S-duality, between the 2 and 3-form theories. They have explained that the S-duality, indeed, is a duality of A and $B+\bar{B}$ models. An evidence for that observation is provided by the BPS counting of black hole states [14]. In the form field theories which we are going to discuss in the present work, one encounters a similar duality of A and $B+\bar{B}$ models but in the case of non-simply connected Calabi-Yau threefolds.

The 7-dimensional manifolds have a specific property which makes them useful to be used in a description of the A and B models in terms of an M-theory. From Abelian duality we know that on every m -dimensional manifold M there is a duality between r -forms and $(m - r - 2)$ -forms [17]. Therefore on a 7-dimensional manifold there is a natural Abelian duality between 2 and 3-form field theories, while in 6 dimensions there does not exist an Abelian duality between 2 and 3-form fields. In this paper we consider the Abelian duality between 2 and 3-form field theories on a trivially fibred $CY_3 \times S^1$. By applying the usual dimensional reduction method we obtain 2 and 3-form field theories on CY_3 which are dual to each other in the sense of [14].

The physical quantities of the form field theories we consider in this paper are by construction elements of $H^{1,1}(CY_3)$, since we are considering a simply connected Calabi-Yau threefold ($h^{1,0} = h^{2,0} = 0$), and $H^3(CY_3) = H^{3,0} \oplus H^{2,1} \oplus H^{1,2} \oplus H^{0,3}$. The theory by construction is independent of the complex structure. But as can be seen from the foregoing Hodge decomposition any representation of the observables amounts to defining a complex structure. Therefore, the physical observables satisfy a set of identities imposed by background independence conditions which ensure the independence of the complex structure [1, 6, 9]. For the 2-form theory, background independence is reasonable if the 2-form field theory is going to be interpreted as the A model string field theory. As we will show the massless sector of the 2-form theory we obtain by dimensional reduction is the theory of Kähler deformation of the Calabi-Yau threefold. To our knowledge such a

theory has been proposed for the first time in [2] where a theory of Kähler deformation was constructed in analogy to the Kodaira-Spencer theory of complex structure deformation, following the mirror symmetry. We show that our proposal for the 2-form theory coincides with the AKS theory of [2] if all terms in the perturbation theory are considered. The 3-form theory studied here is not related to the KS theory since it is the Abelian dual and not the mirror of the 2-form theory and is defined on the same Calabi-Yau threefold and not on its mirror threefold. We further explain this point in terms of a classic/quantum correspondence between the two form field theories.

In this paper we study duality of 2 and 3-form theories which at the classical level correspond to the topological A and B string field theories on the same CY_3 respectively. To this aim we use Abelian duality in 7 dimensions. We explicitly show that the Abelian duality of the 2 and 3-form gravity theories on a trivially fibred $CY_3 \times S^1$ gives, after Kaluza-Klein reduction, an S-duality between 2 and 3-form gravity theories on CY_3 . From a 7 dimensional point of view the Abelian duality gives the position/momentum duality of [14]. Furthermore both in 6 and 7 dimensions the Abelian duality implies a classic/quantum correspondence between the two theories. In fact, the classical regime of the 2-form theory corresponds to the 3-form theory in the $\hbar \rightarrow \infty$ limit. In the classical limit, the 2-form theory reduces to its zero mode sector which is the theory of the Kähler structure deformation. The other extreme is the classical regime of the 3-form field theory which corresponds to the $\hbar \rightarrow \infty$ limit of the 2-form theory. The 3-form theory in this limit reduces to its zero mode sector in which the physical quantities are the elements of $H^3(CY_3)$. It is tempting to anticipate a connection between this theory and the topological B string field theory. But, we notify again that, the 3-form theory should not be expected to be the Kodaira-Spencer theory since one is considering a trivially fibred CY_3 over S^1 .

In a proper gauge we quantize these form gravity theories and describe the Abelian duality as a map between the corresponding annihilation and creation operators. An interesting feature of these theories is the existence of a one parameter family of vacua,

$$|\text{vac.}\rangle_\theta = e^{i\theta a_0} |0\rangle, \quad (2)$$

where a_0 corresponds to the massless modes. The quantum theory of 2-form field theory should be further equipped by suitable holomorphic anomaly equations ensuring the background independence since this is a gravity theory for the A model. On the dual side, quantum background independence for 3-form fields is previously worked out in [6, 7, 9].

The plan of this paper is as follows: in section 2, after a brief review of Abelian duality on general m -dimensional manifolds we study Abelian duality in the case of the 7-dimensional manifold $CY_3 \times S^1$ between 2 and 3-form theories. In a proper gauge the equations of motion of the original field theory reduce to the equations for components of fields along CY_3 . This makes the dimensional reduction of 7-dimensional form field theories to 6 dimensions straightforward. We show that classically the Abelian duality is a position/momentum duality. In the path integral formulation of the Abelian duality we show that it describes a classic/quantum correspondence between the two form field theories. In section 3, we derive 2 and 3-form gravity theories on CY_3 by applying the KK dimensional reduction method to the original 7-dimensional form field theories and reinterpret the classic/quantum correspondence in 6 dimensions. Canonical quantization of the 7-dimensional field theories is studied in section 4. Section 5 is devoted to the connection between the 2-form theory and the theory of Kähler gravity. In section 6 we first construct a method to study the complex structure deformation of a complex manifold. Then we study the quantum background

independence of the massless sector of the 2-form theory. In appendix *A* we derive equations used in section 2. The mathematical background of section 3 is provided in appendix *B*.

2 Abelian duality in 7 dimensions

In this section we derive Abelian duality between 2 and 3-forms in 7 dimensions. For doing this in subsection 2.1 we study Abelian duality of r and $(m - r - 2)$ -forms on an m -dimensional manifold M . Then in subsection 2.2 we use Abelian duality in the 7-dimensional manifold to explain relation between 2 and 3-form theories in 7 dimensions. In subsection 2.3, for the special 7-dimensional manifold $CY_3 \times S^1$ by using gauge fixing condition and after decomposing the components of 2 and 3-forms along CY_3 and S^1 we obtain equations of motion for components of forms along CY_3 and the classical duality between those components. The result is a duality between the 2 and 3-form field theories in 6 dimensions, which is the inheritance of the 7-dimensional Abelian duality.

2.1 Abelian duality

In this subsection we review Abelian duality in m -dimensional Riemannian manifolds M and define form field theories on such manifolds. All of the operators which one encounters in our form theories are shown in the following diagram [17]

$$\begin{array}{ccc} \Omega^r(M) & \xrightarrow{d} & \Omega^{r+1}(M) \\ \downarrow * & & \downarrow * \\ \Omega^{m-r}(M) & \xrightarrow{d^\dagger} & \Omega^{m-r-1}(M) \end{array}$$

where $\Omega^r(M)$ is the set of r -forms on M . If M is a Riemannian manifold, a general r -form η_r satisfies the following equations:

$$d^\dagger \eta_r = (-)^{m(r+1)+1} * d * \eta_r, \quad (3)$$

$$* * \eta_r = (-)^{r(m-r)} \eta_r, \quad (4)$$

where d is the exterior derivative operator and $*$ is the Hodge star operator. There are the same relations with an extra (-1) for Lorentzian manifolds. One can define an inner product of two r -forms η_r and ζ_r on a manifold M ,

$$(\eta_r, \zeta_r) = \int_M \eta_r \wedge * \zeta_r, \quad (5)$$

which is positive definite for Riemannian manifold, $(\varpi_r, \varpi_r) \geq 0$, and the equality is satisfied if and only if $\varpi_r = 0$. Furthermore, $(\eta_r, \zeta_r) = (\zeta_r, \eta_r)$ [18].

For a closed r -form η_r which obeys the condition $d\eta_r = 0$ one can locally assume that $\eta_r = d\zeta_{r-1}$. This follows from the fact that the operator d is nilpotent namely $d^2 = 0$. Also for the nilpotent operator $d^\dagger$ if $d^\dagger \eta_r = 0$ then locally $\eta_r = d^\dagger \zeta_{r+1}$. Therefore if

$$d^\dagger(d\gamma_r) = (-)^{m(r+2)+1} * d * d\gamma_r = 0, \quad (6)$$

in which we have used Eq. (3), then by using Eq. (4), one obtains

$$d(*d\gamma_r) = 0. \quad (7)$$

Thus there exists an $(m - r - 2)$ -form ω_{m-r-2} for which

$$*d\gamma_r = d\omega_{m-r-2}. \quad (8)$$

This relation is known as *Abelian duality*. Considering both the r -form γ_r and the $(m - r - 2)$ -form ω_{m-r-2} as classical fields of two different form field theories, then Eq. (8) is a duality of the corresponding equations of motions.

By using a positive definite inner product we can construct an action for any r -form η_r such that it gives an equation of motion for η_r similar to Eq. (6). This action is defined in the following way:

$$S(\eta) \sim (d\eta_r, d\eta_r) = \int_M d\eta_r \wedge *d\eta_r, \quad (9)$$

for which, using the identity,

$$(d\alpha_r, \beta_r) = (\alpha_r, d^\dagger \beta_r), \quad (10)$$

the equation of motion is given by,

$$d^\dagger d\eta_r = 0. \quad (11)$$

Also, it is obvious that this action is invariant under the following gauge transformation,

$$\eta_r \rightarrow \eta_r + d\theta_{r-1}, \quad (12)$$

where $\theta_{r-1} \in \Omega^{r-1}(M)$. The gauge freedom can be fixed by using the following gauge condition,

$$d^\dagger \eta_r = 0. \quad (13)$$

By using both equation of motion (11) and gauge fixing condition, (13) one obtains that³

$$\Delta \eta_r = 0. \quad (14)$$

Eq. (14) implies that the gauge (13) does not fix the gauge freedom completely since one can still add an arbitrary harmonic form to η .

2.2 7-dimensional Abelian duality

Now, we are ready to study Abelian duality of 2 and 3-form field theories on $\Sigma = CY_3 \times S^1$. Considering $C \in \Omega^3(\Sigma)$ and $K \in \Omega^2(\Sigma)$, we assume that both of these fields obey the following equations of motion⁴

$$d_7^\dagger d_7 C = 0, \quad d_7^\dagger d_7 K = 0, \quad (15)$$

with the Abelian duality relation

$$*_7 d_7 C = d_7 K. \quad (16)$$

To study the quantum version of this classical duality one needs to compare the partition functions of both theories and consequently one should define an action functional for each

³By definition on m -dimensional manifold $\Delta = d^\dagger d + d d^\dagger$.

⁴Subscript of operators d and $*$ refers to the dimensions of manifolds on which the operators are defined.

of them. As we saw in the last subsection, on a compact orientable Riemannian manifold one can define a positive definite action for the 3-form C ,

$$S(C) = \frac{\Lambda}{4\pi} (d_7 C, d_7 C) = \frac{\Lambda}{4\pi} \int_{\Sigma} d_7 C \wedge *_7 d_7 C, \quad (17)$$

which gives Eq. (15) for the 3-form C . The partition function of theory is,

$$Z = \int \mathcal{D}\phi e^{-S(\phi)}. \quad (18)$$

To obtain the Abelian duality, following [19], we introduce new variables such that if we integrate them out from the partition function we get back to the original theory, and if we integrate out the old variables instead of the new ones it produces the dual theory. To this aim we interpret C as a section of a trivial S^1 -bundle $\mathcal{L}_A$ with a connection A , and write an action which includes the covariant derivative $D_{7A}C = d_7C + A$ in the following way,

$$S(C, A) = \frac{\Lambda}{4\pi} \int_{\Sigma} D_{7A}C \wedge *_7 D_{7A}C. \quad (19)$$

There is a simple mechanism for recovering the old action. We introduce a third field K which plays the role of the Lagrange multiplier. We take K to be a section from Σ to S^1 , and write an action,

$$S(C, A, K) = \frac{\Lambda}{4\pi} \int_{\Sigma} D_{7A}C \wedge *_7 D_{7A}C - \frac{i}{2\pi} \int_{\Sigma} K \wedge d_7 A. \quad (20)$$

This action is invariant under the gauge transformation

$$\begin{aligned} A &\rightarrow A + dB, \\ C &\rightarrow C - B, \end{aligned} \quad (21)$$

where B is a 3-form. To see that $S(C, A, K)$ is equivalent to $S(C)$, one can note that the equation of motion for K is $d_7 A = 0$. By imposing this condition and using the $A = 0$ gauge the original theory gets recovered.

The above naive argument can be made exact as follows. Consider the partition function of the whole theory,

$$Z = \frac{1}{\text{vol}(G)} \int \mathcal{D}C \mathcal{D}A \mathcal{D}K \exp \left[-\frac{\Lambda}{4\pi} \int_{\Sigma} D_{7A}C \wedge *_7 D_{7A}C + \frac{i}{2\pi} \int_{\Sigma} K \wedge d_7 A \right], \quad (22)$$

in which G is the gauge group for $\mathcal{L}_A$. By using the delta function,

$$\int \frac{dx}{2\pi} e^{ixy} = \delta(y), \quad (23)$$

the K integral sets dA equal to zero. But we are studying circle-valued forms and we should treat this statement carefully. In general $K = K_{har} + K_{S^1}$, where $K_{har} : \Sigma \rightarrow S^1$ is a circle valued 2-form such that dK_{har} is a harmonic 3-form representative in the cohomology class

of dK , and K_{S^1} is a single valued real function on S^1 . Introducing a basis for harmonic 3-forms λ_j , where $j = 1, \dots, h^3(\Sigma)$,⁵ one can write $dK_{har} = \sum_{j=1}^{h^3(\Sigma)} 2\pi m_j \lambda_j$, in which $m_j \in \mathbb{Z}$. Now one obtains,

$$\begin{aligned} & \int \mathcal{D}K \exp \left[\frac{i}{2\pi} \int_{\Sigma} K \wedge d_7 A \right] \\ &= \int \mathcal{D}K_{S^1} \exp \left[\frac{i}{2\pi} \int_{\Sigma} K_{S^1} \wedge d_7 A \right] \times \sum_{dK_{har} \in H^3(\Sigma, 2\pi\mathbb{Z})} e^{-\frac{i}{2\pi} dK_{har} \wedge A} \\ &= \delta(d_7 A) \prod_j \left(\sum_{m_j \in \mathbb{Z}} e^{-im_j \int_{\Sigma} \lambda_j \wedge A} \right). \end{aligned} \quad (24)$$

The first term shows that A is a flat connection, and among flat connections the gauge equivalence classes $[A]$ are labeled by holonomies, or equivalently by the quantities $\int_{\Sigma} \lambda_j \wedge A$. The remaining part of the integral is gauge invariant therefore one can fix the gauge and integrate over the space of gauge equivalent classes and obtain (omitting the factor $(\text{vol}(G))^{-1}$),

$$\begin{aligned} Z &= \int \mathcal{D}C \mathcal{D}[A] e^{-\frac{\Lambda}{4\pi} \int_{\Sigma} D_7 A C \wedge *7 D_7 A C} \delta(d_7 A) \prod_j \left(\sum_{m_j \in \mathbb{Z}} e^{-im_j \int_{\Sigma} \lambda_j \wedge A} \right) \\ &= \int \mathcal{D}C e^{-\frac{\Lambda}{4\pi} \int_{\Sigma} D_7 A C \wedge *7 D_7 A C} \delta(d_7 A) \delta \left(\int_{\Sigma} \lambda_j \wedge A = 0 \mod 2\pi\mathbb{Z} \right) \\ &= \int \mathcal{D}C e^{-\frac{\Lambda}{4\pi} \int_{\Sigma} d_7 C \wedge *7 d_7 C}. \end{aligned} \quad (25)$$

In the last equality we have used the fact that the delta functions sets the holonomies equal to zero as well as curvature. Thus A is zero modulo gauge transformations, and the new theory is indeed equivalent to the original theory.

Now we want to integrate out C and A to obtain a new theory in terms of the 2-form field K . First, we integrate out C by using the Faddeev-Popov method and then integrate A out by Gaussian method. For the first step, we fix the gauge freedom (21) using the gauge $C = 0$. In this gauge the path integral (22) reduces to,

$$\int \mathcal{D}A \mathcal{D}K \mathcal{D}\bar{c} \mathcal{D}c \exp \left[-\frac{\Lambda}{4\pi} \int_{\Sigma} A \wedge *7 A + \frac{i}{2\pi} \int_{\Sigma} K \wedge d_7 A - \frac{\Lambda}{4\pi} \int_{\Sigma} \bar{c} 1c \right]. \quad (26)$$

The last term arises from gauge fixing by Faddeev-Popov method, and c and $\bar{c}$ are any allowed grassmanian forms on Σ .

For integrating out A , we complete the square. To do this, we introduce new variable $A' = A + \frac{i}{\Lambda} *7 d_7 K$. Then we have,

$$\left(\int \mathcal{D}K e^{-\frac{1}{4\pi\Lambda} \int_{\Sigma} dK \wedge *7 d_7 K} \right) \left(\int \mathcal{D}A' e^{-\frac{\Lambda}{4\pi} \int_{\Sigma} A' \wedge *7 A'} \right) \left(\int \mathcal{D}\bar{c} \mathcal{D}c e^{-\frac{\Lambda}{4\pi} \int_{\Sigma} \bar{c} 1c} \right). \quad (27)$$

The first term is the partition function of the dual theory which is the 2-form theory in 7 dimensions. The last two terms give a topological factor which depends on the number of

⁵ $h^3(\Sigma) = 2 + 2h^{2,1}(CY_3) + h^{1,1}(CY_3)$ [18].

independent harmonic forms in 7-dimensional manifold $CY_3 \times S^1$ (see appendix A for this subtlety). Consequently we have the following equality between the partition functions of two form theories,

$$\int \mathcal{D}C e^{-\frac{\Lambda}{4\pi} \int_{\Sigma} dC \wedge *_{\Sigma} d_7 C} = \mathcal{C}(\Sigma) \int \mathcal{D}K e^{-\frac{1}{4\pi\Lambda} \int_{\Sigma} dK \wedge *_{\Sigma} d_7 K}, \quad (28)$$

It is notable that the coupling Λ in the first theory is mapped to $1/\Lambda$ in the second theory. Therefore there is a classic/quantum correspondence between these form theories. Equation (28) is given in the natural units ($c = \hbar = 1$). Recovering $\hbar$ in Eq. (28) one obtains $Z(\phi) = \int \mathcal{D}\phi e^{-\frac{1}{\hbar} S(\phi)}$. Thus, in 3-form (2-form) theory $1/\Lambda$ (Λ) acts as $\hbar$ and we can study classical and quantum regimes of each theory in the limit of Λ tending to 0 or ∞ .

2.3 Equation of motion and gauge fixing

The equation of motion for 3-form C given by the action (17) is,

$$d_7^\dagger d_7 C = 0. \quad (29)$$

After the decomposition $C = \gamma + dt \wedge \alpha$ where γ is a 3-form along CY_3 and $dt \wedge \alpha$ is a 1-form along S^1 and a 2-form along CY_3 , and also the decomposition of the operator d_7 into its components along CY_3 and S^1 , $d_7 = d_{CY} + dt \wedge d/dt$, the equation of motion (29) reduces to,⁶

$$d_{CY}^\dagger d_{CY} \gamma - \ddot{\gamma} - d_{CY} \dot{\alpha} = 0, \quad (30)$$

$$d_{CY}^\dagger d_{CY} \alpha + d_{CY}^\dagger \dot{\gamma} = 0, \quad (31)$$

where dot “.” stands for a derivation with respect to t which parameterizes the S^1 . To obtain this results one needs the relations between operators d and $*$ in the 7-dimensional manifold Σ and the 6-dimensional manifold CY_3 which can be found in appendix B.

The action (17) is invariant under the gauge transformation $C \rightarrow C + d_7 \rho$, where ρ is a 2-form along Σ . We can fix this gauge freedom by $d_7^\dagger C = 0$ gauge. In terms of γ and α , the gauge transformation is,

$$\gamma \rightarrow \gamma + d\rho_1, \quad \alpha \rightarrow \alpha + \dot{\rho}_1 + d\rho_2, \quad (32)$$

in which $\rho = \rho_1 + \rho_2 \wedge dt$ and the gauge fixing condition is equivalent to,

$$d_{CY}^\dagger \gamma + \dot{\alpha} = 0, \quad (33)$$

$$d_{CY}^\dagger \alpha = 0. \quad (34)$$

By operating d_{CY} and d/dt on the first equation and inserting the result into the equations of motion (30,31), one obtains,

$$\Delta_{CY} \gamma - \ddot{\gamma} = 0, \quad (35)$$

$$\Delta_{CY} \alpha - \ddot{\alpha} = 0. \quad (36)$$

⁶We can obtain these equations of motion by putting decomposition of C in action(17) and using Euler-Lagrange equation.

Similar calculations can be done in the case of the 2-form theory in 7 dimensions. From Eq. (28) one verifies that the action for K is,

$$S(K) = \frac{1}{4\pi\Lambda} \int_{\Sigma} d_7 K \wedge *_7 d_7 K. \quad (37)$$

After the decomposition $K = \omega + dt \wedge \beta$, where ω and β are 2-form and 1-form along CY_3 respectively, the equation of motion for K , i.e. $d_7^\dagger d_7 K = 0$, reduces to,

$$d_{CY}^\dagger d_{CY} \omega - \ddot{\omega} + d_{CY} \dot{\beta} = 0, \quad (38)$$

$$d_{CY}^\dagger d_{CY} \beta - d_{CY}^\dagger \dot{\omega} = 0. \quad (39)$$

The gauge symmetry of the action for 2-form K is,

$$K \rightarrow K + d\theta, \quad (40)$$

where θ is an 1-form along Σ . The gauge fixing condition $d_7^\dagger K = 0$ gives,

$$d_{CY}^\dagger \omega - \dot{\beta} = 0, \quad (41)$$

$$d_{CY}^\dagger \beta = 0, \quad (42)$$

which can be used to simplify the equations of motion (38,39) as follows,

$$\Delta_{CY} \omega - \ddot{\omega} = 0, \quad (43)$$

$$\Delta_{CY} \beta - \ddot{\beta} = 0. \quad (44)$$

The classical duality (16) between the 3-form C and 2 form K results in the following duality relations,

$$*_CY d_{CY} \gamma = \dot{\omega} + d_{CY} \beta, \quad (45)$$

$$*_CY (\dot{\gamma} + d_{CY} \alpha) = d_{CY} \omega. \quad (46)$$

In the next sections we will argue that these relations give us an interpretation of the Abelian duality as the position/momentum duality [14].

The 3-form action (17) in terms of the γ and α components is,⁷

$$S(\gamma, \alpha) = \frac{\Lambda}{4\pi} \int_{S^1} dt \{(\dot{\gamma}, \dot{\gamma}) + (d\gamma, d\gamma) + 2(\dot{\gamma}, d\alpha) + (d\alpha, d\alpha)\}. \quad (47)$$

Since we are interested in form field theories on CY_3 , we use the gauge freedom (32) to set,

$$\alpha = 0. \quad (48)$$

In this gauge the action (47) simplifies to,

$$S(\gamma) = \frac{\Lambda}{4\pi} \int_{S^1} dt \{(\dot{\gamma}, \dot{\gamma}) + (d\gamma, d\gamma)\}. \quad (49)$$

⁷Henceforth, all operators and the corresponding inner products are defined on CY_3 and we omit the subscript CY .

One should note that there is still a gauge freedom,

$$\gamma \rightarrow \gamma + d\tilde{\rho}_1, \quad (50)$$

where $\tilde{\rho}_1$ is an arbitrary 2-form independent of t ,

$$\tilde{\rho}_1 \in \Omega^2(CY_3). \quad (51)$$

By the same argument one can get rid of the 1-form β in the 2-form theory to obtain $S(\omega) = \frac{1}{4\pi\Lambda} \int_{S^1} dt \{(\dot{\omega}, \dot{\omega}) + (d\omega, d\omega)\}$. In these gauges the classical dualities (45) and (46) reduce to,

$$*d\gamma = \dot{\omega}, \quad (52)$$

$$*\dot{\gamma} = d\omega. \quad (53)$$

3 Reduction to Calabi-Yau threefold

In this section we study the reduction of 2 and 3-form theories in 7 dimensions to form gravity theories on 6-dimensional manifold CY_3 by using Kaluza-Klein method. We assume that R is the radius of S^1 over which the CY_3 is fibred trivially. In this way we can expand all forms,

$$\gamma = \sum_{n \in \mathbb{Z}} \gamma_n e^{i2\pi n t/R}, \quad (54)$$

$$\omega = \sum_{n \in \mathbb{Z}} \omega_n e^{i2\pi n t/R}, \quad (55)$$

where γ_n and ω_n are defined on CY_3 and their dependence on the extra dimension t along S^1 is given in the phase $e^{i2\pi n t/R}$. Thus both actions on the CY_3 reduce to,

$$S(\gamma_n) = \frac{\Lambda}{4\pi} \sum_{n \in \mathbb{Z}} \left\{ \frac{4\pi^2 n^2}{R^2} (\gamma_n, \gamma_{-n}) + (d\gamma_n, d\gamma_{-n}) \right\}, \quad (56)$$

$$S(\omega_n) = \frac{1}{4\pi\Lambda} \sum_{n \in \mathbb{Z}} \left\{ \frac{4\pi^2 n^2}{R^2} (\omega_n, \omega_{-n}) + (d\omega_n, d\omega_{-n}) \right\}. \quad (57)$$

To obtain the above results the following identity is used,

$$\int_{S^1} dt e^{i2\pi(m+n)t/R} = \delta_{m+n,0} \quad (58)$$

In addition, all of the equations of motion on CY_3 in terms of KK modes simplify to,

$$d^\dagger d\gamma_n = -\frac{4\pi^2 n^2}{R^2} \gamma_n, \quad (59)$$

$$d^\dagger d\omega_n = -\frac{4\pi^2 n^2}{R^2} \omega_n. \quad (60)$$

We see that coefficient the $\frac{4\pi^2 n^2}{R^2}$ in both theories gives the KK mass squared of the n -th mode.

There is a subtlety in the partition functions of these theories as can be seen from Eqs.(56,57). In the classical limit of the 3-form (2-form) theory $\Lambda \rightarrow \infty$ ($1/\Lambda \rightarrow \infty$), which corresponds to $\hbar \rightarrow 0$, the partition function gets localized around the classical trajectory. By using equations of motion (59,60) one can obtain the following inner product for the KK modes on the Calabi-Yau threefold

$$\begin{aligned}(d\gamma_n, d\gamma_n) &= -\frac{4\pi^2 n^2}{R^2} (\gamma_n, \gamma_n), \\ (d\omega_n, d\omega_n) &= -\frac{4\pi^2 n^2}{R^2} (\omega_n, \omega_n).\end{aligned}\tag{61}$$

Recalling that for Riemannian manifolds the inner product is positive definite, one verifies that the solution of the equations of motion is $\gamma_n = 0 = \omega_n$ for $n \neq 0$. Consequently in the classical limit each theory reduces to its zero mode sector. Furthermore, one easily verifies the classic/quantum correspondence between the two form field theories is a correspondence between the massless sector of one theory and the full spectrum of the another one. In section 5 we will show that the massless sector of the 2-form theory gives the theory of Kähler gravity on the CY_3 . Thus the classic/quantum correspondence implies why the massless sector of the 3-form field theory is not the Kodaira-Spencer theory which is believed to correspond to the topological B string field theory.

4 Canonical quantization and position/momentum duality

Now we are able to do canonical quantization. Recall that the actions of 3 and 2-form theories in 7 dimensions respectively are,

$$S(\gamma) = \frac{\Lambda}{4\pi} \int_{S^1} dt \{(\dot{\gamma}, \dot{\gamma}) + (d\gamma, d\gamma)\},\tag{62}$$

$$S(\omega) = \frac{1}{4\pi\Lambda} \int_{S^1} dt \{(\dot{\omega}, \dot{\omega}) + (d\omega, d\omega)\}.\tag{63}$$

The momenta conjugate to γ and ω are respectively,⁸

$$P_\gamma = \frac{\Lambda}{2\pi} \dot{\gamma},\tag{65}$$

$$P_\omega = \frac{1}{2\pi\Lambda} \dot{\omega}.\tag{66}$$

The Poisson-bracket is given as usual e.g.

$$\{\gamma, P_\gamma\} = 1,\tag{67}$$

⁸Here we have used the following definition,

$$\frac{\partial(\psi, \chi)}{\partial\psi} = \frac{\partial(\chi, \psi)}{\partial\psi} = \chi,\tag{64}$$

which means that the derivation is defined with respect to the *product* $\wedge*$.

which results in,

$$\{\gamma_n, \gamma_m\} = \frac{i}{n} \frac{R}{\Lambda} \delta_{n,-m}, \quad n, m \neq 0, \quad (68)$$

$$\{\gamma_0, \gamma_n\} = 0, \quad n \in \mathbb{Z}. \quad (69)$$

One can define the creation and annihilation operators as usual

$$a_n = \sqrt{n} \gamma_n, \quad n > 0, \quad (70)$$

$$a_{-n}^\dagger = \sqrt{-n} \gamma_n, \quad n < 0, \quad (71)$$

$$a_0 = a_0^\dagger = \gamma_0, \quad (72)$$

satisfying the algebra

$$[a_n, a_m^\dagger] = \frac{R}{\Lambda} \delta_{n,m}, \quad n, m \neq 0, \quad (73)$$

$$[a_0, a_n] = [a_0, a_n^\dagger] = 0, \quad n \in \mathbb{Z}. \quad (74)$$

The vacuum state for the 3-form theory $|0\rangle$ is defined to be the state annihilated by a_n . There is a subtlety at this point. Since a_0 commutes with the massive operators $a_n^\dagger$, one verifies that this operator takes one vacuum state into a nearby equivalent one [11]. Therefore one can define a one parameter family of vacua such that,

$$|\text{vac.}\rangle_\theta = e^{i\theta a_0} |0\rangle. \quad (75)$$

We can quantize the 2-form theory (63) in the same way and derive the associated vacua family. The difference between these two form field theories is due to the exponent of Λ in the commutation algebra. In the 2-form theory the commutation relation is given by,

$$[b_n, b_m^\dagger] = R\Lambda \delta_{n,m}, \quad n, m \neq 0, \quad (76)$$

$$[b_0, b_n] = [b_0, b_n^\dagger] = 0, \quad n \in \mathbb{Z}, \quad (77)$$

where

$$b_n = \sqrt{n} \omega_n, \quad n > 0, \quad (78)$$

$$b_{-n}^\dagger = \sqrt{-n} \omega_n, \quad n < 0, \quad (79)$$

$$b_0 = b_0^\dagger = \omega_0. \quad (80)$$

Consequently the classical limit of one theory $\hbar \rightarrow 0$ (e.g. for 3-form theory it means $\Lambda \rightarrow \infty$) corresponds to the quantum extreme of the another theory. This is the classic/quantum correspondence of 2 and 3-form theories in CY_3 induced by the Abelian duality.

4.1 S-duality on CY_3

Now we can understand the duality between the two form gravity theories. The duality is two-edged indeed. Firstly, there is a classic/quantum correspondence between two theories. Secondly, from the classical duality relations (52,53) and definition of momentum (65,66) we obtain the interesting relations,

$$*d\gamma = 2\pi\Lambda P_\omega, \quad (81)$$

$$*P_\gamma = \frac{\Lambda}{2\pi} d\omega, \quad (82)$$

which can be interpreted as position/momentum or electric/magnetic duality in 7-dimensional form theories[14]. In the 6 dimensional manifold, this relations in terms of KK modes reduce to,

$$*d\gamma_n = i\frac{2\pi n}{R}\omega_n, \quad (83)$$

$$*d\omega_n = i\frac{R}{2\pi n}\gamma_n. \quad (84)$$

There is a subtlety in these equations. We see that because of the coefficient n , the zero modes of the theories do not play any role in the above duality relations. Zero modes play a significant role in the classical/quantum correspondence. Here we recall our discussion on the classic/quantum correspondence at the end of section 3. In the classical limit, the 3-form field theory for example, is given by its massless sector which corresponds to the 2-form theory in its quantum extreme with both massless and massive excitations.

5 Theory of Kähler gravity

Let us see what is the relation between the action,

$$S(\omega) \sim (d\omega, d\omega), \quad (85)$$

and the theory of Kähler gravity. It is known that the observables of the A-model are elements of $H^{(p,q)}$. For $p = q = 1$ one verifies a correspondence between the A-model and the theory of Kähler transformations.

In physical superstring theories we have $N = (2, 2)$ supersymmetry. After the twist to the topological string theories there remain only two supercharges. In the B string field theory the corresponding nilpotent operators are ∂ and $\bar{\partial}$. A deformation in the complex structure amounts to deformation of $\bar{\partial}$ given by,

$$\bar{\partial} \rightarrow \bar{\partial} + A.\partial, \quad (86)$$

where $A \in H^{0,1}(TM)$. Requiring that the deformed operator is nilpotent one obtains the Kodaira-Spencer equation[3],

$$\bar{\partial}A' + \frac{1}{2}\partial(A \wedge A)' = 0, \quad (87)$$

where $A' = A.\Omega$ in which Ω is the Calabi-Yau three form.

In the A model the nilpotent operators are d and $d^{c\dagger}$.⁹ Operator d is nilpotent without any dependence on the Kähler and/or complex structure. Given a Kähler structure ω_0 the $d^{c\dagger}$ operator is defined as follows,

$$d^{c\dagger} = [d, \Lambda_0], \quad (88)$$

where Λ_0 is the pull back of multiplication by ω_0 [20]. As we see, the definition of the operators d and $d^{c\dagger}$ is independent of the complex structure, but for $d^{c\dagger}$ it depends on the

⁹By definition we have:

$$d = \partial + \bar{\partial}, \quad d^c = \bar{\partial} - \partial.$$

One can define d without determining complex structure but define $d^\dagger$ one should choose a complex structure.

Kähler structure as is expected for the A string field theory. By a Kähler transformation one means a deformation of the Kähler form ω_0 ,

$$\omega_0 \rightarrow \omega = \omega_0 + K, \quad (89)$$

where K is a 2-form, leaving the operator $D^{c\dagger} = d^{c\dagger} + [d, \delta\Lambda]$ be nilpotent. Again $\Lambda = \Lambda_0 + \delta\Lambda$ is the pull back of the multiplication by ω . Given a complex structure, Λ in components is given by the following identities,

$$\Lambda^{i\bar{j}} \omega_{i\bar{k}} = \delta_{\bar{k}}^{\bar{j}}, \quad (90)$$

$$\Lambda^{i\bar{j}} \omega_{k\bar{j}} = \delta_k^i. \quad (91)$$

For a general Kähler form ω it is straightforward to verify that $d^{c\dagger} = [d, \Lambda]$ is nilpotent if and only if,

$$\Lambda_{,k}^{l\bar{j}} \Lambda^{k\bar{m}} = \Lambda_{,k}^{l\bar{m}} \Lambda^{k\bar{j}}, \quad (92)$$

and with the same symmetry for the holomorphic indices. Here by “ $,k$ ” one means derivation with respect to the holomorphic coordinate z^k . The identity (92) is satisfied *iff* the inverse of Λ i.e. the two form ω is closed, namely,

$$\omega_{i\bar{j},k} = \omega_{k\bar{j},i}, \quad \omega_{i\bar{j},\bar{k}} = \omega_{i\bar{k},\bar{j}}. \quad (93)$$

Thus the Kähler transformation is given by closed 2-forms K in (89). An even dimensional Riemannian manifold can always be equipped with a complex structure for which the action functional,

$$S \sim (d\omega, d\omega), \quad (94)$$

is positive definite implying the desired condition,

$$d\omega = 0, \quad (95)$$

by the classical equation of motion.

In summary the massless sector of the 2-form theory studied in the previous sections corresponds to the A string field theory.

Here we slightly digress to discuss the difference between our method and that of [2] where the theory of Kähler gravity is considered for the first time. The motivation in [2] is as obtain a theory of Kähler transformation to be the mirror of the Kodaira-Spencer theory of deformation of complex structure moduli. The so called AKS theory of Kähler transformations is given by requiring that the operator

$$D = d + [d^{c\dagger}, K], \quad (96)$$

be nilpotent. On a manifold the operator d can be defined with no reference to the complex structure. Given a Kähler structure by ω and the corresponding pull back operator Λ one can define the operator $d^{c\dagger}$ by Eq. (88) which itself depends on ω . Thus the AKS condition,

$$D^2 = 0, \quad (97)$$

is well defined only by perturbation as far as $d^{c\dagger}$ in Eq. (96) is defined in terms of d and $\omega = \omega_0 + K$. In our approach one uses the fact that the operator d is defined independent

of the Kähler structure and the equation of motion of the A string field theory is obtained from the condition,

$$(d^{\dagger})^2 = [d, \Lambda]^2 = 0. \quad (98)$$

Thus our proposal is equivalent to the AKS conjecture if all terms in the perturbation are considered.

There are some subtleties in defining the action (94). The action S in Eq. (94),

$$S \sim \int dK \wedge *dK, \quad (99)$$

is defined perturbatively since the Hodge $*$ operation is defined in terms of Λ which is known in terms of the deformation K only via the following equation,

$$\Lambda^{i\bar{j}} = \Lambda_0^{i\bar{j}} - \Lambda_0^{i\bar{k}} K_{k\bar{k}} \Lambda_0^{k\bar{j}} + \dots. \quad (100)$$

Up to $\mathcal{O}(K^5)$ terms the action is given by as follows,

$$\begin{aligned} S(K) &\sim \int dV (K_{i\bar{j},\bar{k}} K_{j\bar{i},k} + K_{j\bar{i},k} K_{i\bar{j},\bar{k}}) \Lambda^{i\bar{i}} (\Lambda^{j\bar{j}} \Lambda^{k\bar{k}} - j \leftrightarrow k) \\ &= \int dV (K_{i\bar{j},\bar{k}} K_{j\bar{i},k} + K_{j\bar{i},k} K_{i\bar{j},\bar{k}}) \times \\ &\quad \left\{ \left(\Lambda_0^{i\bar{i}} \Lambda_0^{j\bar{j}} \Lambda_0^{k\bar{k}} \right) \right. \\ &\quad - \left(\Lambda_0^{i\bar{l}} K_{l\bar{l}} \Lambda_0^{l\bar{i}} \Lambda_0^{j\bar{j}} \Lambda_0^{k\bar{k}} + \Lambda_0^{i\bar{i}} \Lambda_0^{j\bar{l}} K_{l\bar{l}} \Lambda_0^{l\bar{j}} \Lambda_0^{k\bar{k}} + \Lambda_0^{i\bar{i}} \Lambda_0^{j\bar{j}} \Lambda_0^{k\bar{l}} K_{l\bar{l}} \Lambda_0^{l\bar{k}} \right) \\ &\quad + \left(\Lambda_0^{i\bar{l}} K_{l\bar{l}} \Lambda_0^{l\bar{i}} \Lambda_0^{j\bar{m}} K_{m\bar{m}} \Lambda_0^{m\bar{j}} \Lambda_0^{k\bar{k}} + \Lambda_0^{i\bar{l}} K_{l\bar{l}} \Lambda_0^{l\bar{i}} \Lambda_0^{j\bar{j}} \Lambda_0^{k\bar{m}} K_{m\bar{m}} \Lambda_0^{m\bar{k}} \right. \\ &\quad + \Lambda_0^{i\bar{i}} \Lambda_0^{j\bar{l}} K_{l\bar{l}} \Lambda_0^{l\bar{j}} \Lambda_0^{k\bar{m}} K_{m\bar{m}} \Lambda_0^{m\bar{k}} + \Lambda_0^{i\bar{l}} K_{l\bar{l}} \Lambda_0^{l\bar{m}} K_{m\bar{m}} \Lambda_0^{m\bar{i}} \Lambda_0^{j\bar{j}} \Lambda_0^{k\bar{k}} \\ &\quad + \Lambda_0^{i\bar{i}} \Lambda_0^{j\bar{l}} K_{l\bar{l}} \Lambda_0^{l\bar{m}} K_{m\bar{m}} \Lambda_0^{m\bar{j}} \Lambda_0^{k\bar{k}} + \Lambda_0^{i\bar{i}} \Lambda_0^{j\bar{j}} \Lambda_0^{k\bar{l}} K_{l\bar{l}} \Lambda_0^{l\bar{m}} K_{m\bar{m}} \Lambda_0^{m\bar{k}} \left. \right) \\ &\quad \left. + \mathcal{O}(K^5) - (j \leftrightarrow k) \right\}. \end{aligned} \quad (101)$$

The gauge condition,

$$d^{\dagger} K = 0, \quad (102)$$

implies that, e.g.

$$\Lambda_0^{i\bar{j}} K_{i\bar{l},\bar{j}} + \mathcal{O}(K^2) = 0. \quad (103)$$

Thus K in this gauge is given by *transverse* modes, a well known notion in the ordinary quantum gravity.

The action (99) is appeared to depend on the complex structure though the classical equation $d\omega = 0$ was obtained with no reference to the complex structure. For simply connected manifolds, the independence of the action from the complex structure can be verified as follows. Any variation of the complex structure deforms the operator $\bar{\partial}$ by $A \in H^{0,1}(TM)$ which is the Kodaira-Spencer field [3]. Since $A.K$ is a (0,2)-form, for a

simply connected manifold it is vanishing identically,

$$\begin{aligned}
A.K &= (A.K)_{i\bar{j}} d\bar{z}^i \wedge d\bar{z}^j \\
&= d\eta \\
&= d(\eta_i dz^i + \eta_{\bar{i}} d\bar{z}^{\bar{i}}) \\
&= d(\eta_i dz^i) + d(\eta_{\bar{i}} d\bar{z}^{\bar{i}}) \\
&= d(d(\phi^{(1)})) + d(d(\phi^{(2)})) \\
&= 0,
\end{aligned} \tag{104}$$

where η and $\phi^{(i)}$ are 1 and 0-forms on simply connected manifold respectively. In the 2nd and 5th lines we used the fact that for a simply connected CY_3 , $h^{0,2} = h^{0,1} = 0$. Consequently the theory is classically protected against complex structure deformation. In the next section we study the quantum background independence of this theory.

Before closing this section we address the modification in the action (94) for non-simply connected manifolds if it is used to describe the A string field theory. The A model should be independent of the complex structure moduli. Thus one should add the condition

$$A.K = 0, \tag{105}$$

to the action using the Lagrange multipliers α_i ,

$$S = (d\omega, d\omega) + \left\{ \alpha_i \int_{C_i} A.K + h.c. \right\}, \tag{106}$$

where C_i 's are $(0, 2)$ cycles. The second term induces a coupling between the A-model and B+ $\bar{B}$ model.¹⁰ For a Calabi-Yau manifold the constraint (105) can be equivalently given by

$$A' \wedge K = 0. \tag{107}$$

Thus the modified action can be given by

$$S = (d\omega, d\omega) + \left\{ \int_{CY_3} \lambda \wedge A' \wedge K + h.c. \right\}, \quad \lambda \in H^{0,1}. \tag{108}$$

We postpone further investigation of this theory to future works.

6 Quantum background independence

The observables of the A string field theory should be independent of the complex structure. In this section at first we construct the necessary tools to study the theory of complex structure deformation, then we use these tools to study the quantum background independence of $H^2(CY_3)$ and $H^3(CY_3)$. The quantum background independence of $H^3(CY_3)$ is worked out previously and is known to result in the holomorphic anomaly equations [5, 6, 7, 9].

¹⁰In [14], in a different approach to form field theories on CY_3 , it is argued that on a Calabi-Yau threefold, the A model is dual to the B+ $\bar{B}$ model.

6.1 The structure of connections

In this subsection we study the connections corresponding to the deformation of the complex structure on a complex manifolds. It is known that the number of moduli complex structure is $h^{2,1}$. An infinitesimal deformation of complex structure makes the holomorphic differentials dz^i on the complex manifold to be mixed with the anti-holomorphic differentials [21],

$$\frac{\partial}{\partial t_M} dz^i = \nu_{jM}^i dz^j + \mu_{jM}^i d\bar{z}^{\bar{j}}, \quad (109)$$

where t_M are variables of complex structure moduli space and M counts the number of complex structure moduli, $M = 1, \dots, h^{2,1}$. Fundamental duality relation between basis of forms and basis of vectors,

$$\begin{aligned} \langle dz^i, \bar{\partial}_{\bar{j}} \rangle &= \langle d\bar{z}^{\bar{i}}, \partial_j \rangle = 0, \\ \langle dz^i, \partial_j \rangle &= \delta_j^i, \\ \langle d\bar{z}^{\bar{i}}, \bar{\partial}_{\bar{j}} \rangle &= \delta_{\bar{j}}^{\bar{i}}, \end{aligned} \quad (110)$$

should be satisfied after the deformation. Therefore we obtain the following relations for those basis,

$$\begin{aligned} \frac{\partial}{\partial t_M} dz^i &= \nu_{jM}^i dz^j + \mu_{jM}^i d\bar{z}^{\bar{j}}, \\ \frac{\partial}{\partial t_M} \partial_i &= -\nu_{iM}^j \partial_j + \epsilon_{iM}^{\bar{j}} \bar{\partial}_{\bar{j}}, \\ \frac{\partial}{\partial t_M} d\bar{z}^{\bar{i}} &= \xi_{jM}^{\bar{i}} d\bar{z}^{\bar{j}} - \epsilon_{jM}^i dz^j, \\ \frac{\partial}{\partial t_M} \bar{\partial}_{\bar{i}} &= -\xi_{iM}^{\bar{j}} \bar{\partial}_{\bar{j}} - \mu_{iM}^{\bar{j}} \bar{\partial}_{\bar{j}}. \end{aligned} \quad (111)$$

Thus we have four sets of independent connections μ_M , ν_M , ϵ_M , and ξ_M of complex structure deformation on a complex manifold. In the next subsection we show on a CY_3 manifold only two sets of these connections remain and two other sets are vanishing identically.

6.2 Complex structure independence in form field theories

In the 3-form gravity theory, which is defined on a Calabi-Yau threefold manifold, real closed 3-form γ_0 can be decomposed in the basis consisting of the Calabi-Yau 3-form $\Omega \in H^{3,0}$,¹¹ its covariant derivatives with respect to the moduli directions t_M , $\nabla_M \Omega \in H^{2,1}$,¹² and their complex conjugates,

$$\gamma_0 = \lambda^{-1} \Omega + x^M \nabla_M \Omega + \bar{x}^{\bar{M}} \bar{\nabla}_{\bar{M}} \bar{\Omega} + \bar{\lambda}^{-1} \bar{\Omega}. \quad (112)$$

¹¹For Calabi-Yau threefold we have $h^{0,0} = h^{3,0} = h^{0,3} = h^{3,3} = 1$

¹² $H^3(CY_3)$ forms a bundle over the moduli space of the complex structures of the CY_3 . The holomorphic 3-form defines a line sub-bundle $\mathcal{L}$ and on this sub-bundle one can define a connection $\nabla_M = \partial_M + \partial_M K$ where K is the Kähler potential defined as follows,

$$K = -\ln i \int \bar{\Omega} \wedge \Omega.$$

Locally the Calabi-Yau 3-form takes the form,

$$\Omega = \frac{1}{3!} h(z) \varepsilon_{ijk} dz^i \wedge dz^j \wedge dz^k, \quad (113)$$

and

$$\frac{\partial}{\partial t_M} \Omega = \frac{1}{3!} h(z)_{,M} \varepsilon_{ijk} dz^i \wedge dz^j \wedge dz^k + \frac{1}{2!} h(z) \varepsilon_{ijk} \frac{\partial}{\partial t_M} (dz^i) \wedge dz^j \wedge dz^k. \quad (114)$$

By expanding this in terms of μ_{jM}^i and ν_{jM}^i and using Eqs. (111) one verifies that μ_{jM}^i and ν_{jM}^i are generators of bidegree (2,1) and (3,0)-forms respectively.

Since on a Calabi-Yau threefold $\partial_M \bar{\Omega} = 0$ and Ω is a nowhere vanishing holomorphic 3-form, by expanding $\partial_M \bar{\Omega}$ one can verify that $\xi_{bM}^{\bar{a}} = 0 = \epsilon_{bM}^a$ and Eqs.(111) simplify to

$$\begin{aligned} \frac{\partial}{\partial t_M} dz^i &= \nu_{jM}^i dz^j + \mu_{jM}^i d\bar{z}^{\bar{j}}, \\ \frac{\partial}{\partial t_M} \partial_i &= -\nu_{iM}^j \partial_j, \\ \frac{\partial}{\partial t_M} d\bar{z}^{\bar{i}} &= 0, \\ \frac{\partial}{\partial t_M} \bar{\partial}_{\bar{i}} &= -\mu_{iM}^{\bar{j}} \bar{\partial}_{\bar{j}}. \end{aligned} \quad (115)$$

Consequently all of the complex structure deformation on Calabi-Yau threefold are given by μ_{bM}^a and ν_{bM}^a .

In 3-form theory it is shown that the holomorphic anomaly equations depend only on the μ_{bM}^a connections [4] which can be equivalently obtained by imposing the background independence condition on $H^3(CY_3)$ [6, 9].

For the 2-form gravity theory one deals with a 2-form ω_0 on the CY_3 . The 2-form ω_0 is given in terms of a basis of $H^2(CY_3)$ as follows,

$$\omega_0 = x_I \omega^I(z), \quad (116)$$

where I runs over the basis of $H^2(CY_3)$. If a compact simply connected CY_3 manifold is considered, since for any suitable complex structure we have $h^{2,0} = 0$ any basis for $H^2(CY_3)$ is a basis for the Kähler forms. For a given complex structure $\omega^I(z) = \omega^I(z)_{a\bar{b}} dz^a \wedge d\bar{z}^{\bar{b}}$ for $I = 1, \dots, h^{1,1}$. Thus one obtains,

$$\partial_M \omega^I = (\omega_{i\bar{j}}^I)_{,M} dz^i \wedge d\bar{z}^{\bar{j}} + \omega_{i\bar{j}}^I \partial_M (dz^i) \wedge d\bar{z}^{\bar{j}}. \quad (117)$$

By expanding this formula in terms of μ_{jM}^i and ν_{jM}^i and noting that $h^{2,0}(CY_3) = 0$ one obtains that μ_{jM}^i is a null operator on the H^2 i.e.

$$\mu_{iM}^j \omega_{j\bar{i}}^I d\bar{z}^{\bar{i}} \wedge d\bar{z}^{\bar{j}} = 0. \quad (118)$$

By imposing the background (i.e. complex structure) independence condition on the symplectic 2-form ω_0 , one obtains,

$$[\nu_{Mi}^j \omega_{j\bar{i}}^I + (\omega_{i\bar{i}}^I)_{,M}] dz^i \wedge d\bar{z}^{\bar{i}} = 0. \quad (119)$$

Thus, ν_{Mi}^j are generators of transformation along the complex structure moduli directions for $H^2(CY_3)$.

A Abelian duality in 7 dimensions

At the end of section 2 we saw that a coefficient multiplies the partition functions of 2-form theory in 7 dimensions. In this appendix we show that this coefficient is a topological invariant which only depends on the Hodge numbers of CY_3 . For deriving that coefficient one should calculate two terms; the Gaussian integral over A' , $\int \mathcal{D}A' e^{-\frac{\Lambda}{4\pi} \int_{\Sigma} A' \wedge *_{\Sigma} A'}$, and the grassmanian integral over c and $\bar{c}$, $\int \mathcal{D}\bar{c} \mathcal{D}c e^{-\frac{\Lambda}{4\pi} \int_{\Sigma} \bar{c} c}$. For the first one, we note that A' is a 4-form and the Hodge star operator is a linear map. Thus this term gives a factor of $\int dy e^{-(\Lambda/4\pi)y} = \sqrt{\frac{\pi}{\Lambda/4\pi}}$ for each 4-form.¹³

For the second term it should be noted that the path integral over the ghost fields run over the set of grassmanian r -forms and $(7-r)$ -forms where $r = 0, \dots, 7$ (note that either the r or $(7-r)$ -form is an even form). For each set of ghost fields one obtains $\int \mathcal{D}\bar{c} \mathcal{D}c e^{-\frac{\Lambda}{4\pi} \int_{\Sigma} \bar{c} c} = \Lambda/4\pi$. Thus the overall factor is

$$\mathcal{C}(\Sigma) = (2\pi)^{1+h^{2,1}+h^{1,1}/2} (\Lambda/2\pi)^{4+h^{2,1}+3h^{1,1}/2}, \quad (120)$$

where we have used the following identities for the Hodge numbers of $\Sigma = CY_3 \times S^1$ [18],

$$\begin{aligned} b^0(\Sigma) &= b^7(\Sigma) = 1, \\ b^1(\Sigma) &= b^6(\Sigma) = 2, \\ b^2(\Sigma) &= b^5(\Sigma) = h^{1,1}(CY_3), \\ b^3(\Sigma) &= b^4(\Sigma) = 2h^{2,1}(CY_3) + h^{1,1}(CY_3) + 2. \end{aligned} \quad (121)$$

B Relation between the operators d and $*$ in m and $m+1$ dimensions

In this appendix we deal with some relations between the operators in m and $m+1$ dimensions. In section 2 we used two elementary operators, namely the exterior derivative d and the Hodge star $*$ in 6 and 7 dimensions.

Operator d is a map $d : \Omega^r(M) \rightarrow \Omega^{r+1}(M)$, where $\Omega^r(M)$ are r -forms of manifold M , whose action on a r -form,

$$\eta_r = \frac{1}{r!} \eta_{\mu_1 \dots \mu_r} dx^{\mu_1} \wedge \dots \wedge dx^{\mu_r}, \quad (122)$$

is defined by,

$$d\eta_r = \frac{1}{r!} \left(\frac{\partial}{\partial x^\nu} \eta_{\mu_1 \dots \mu_r} \right) dx^\nu \wedge dx^{\mu_1} \wedge \dots \wedge dx^{\mu_r}. \quad (123)$$

To obtain the relation between exterior derivative in m and $m+1$ dimensional manifolds, one should note that for our purpose $(m+1)$ -th direction in $(m+1)$ -dimensional manifold is perpendicular to other directions. Therefore it is obvious that,

$$d_{m+1} = d_m + dt \wedge \frac{\partial}{\partial t}, \quad (124)$$

where subscripts m and $m+1$ indicate the dimension of manifold and t is the coordinate of $(m+1)$ -th direction.

¹³ y is a normalization real variable and is defined such that $\int \mathcal{D}A' e^{-\int_{\Sigma} A' \wedge * A'} = \int dy e^{-y}$ [22].

For Hodge star operator there are some subtleties. Operator $*$ is a linear map $\Omega^r(M) \rightarrow \Omega^{m-r}(M)$, where m is the dimension of manifold M . The operation of the Hodge $*$ on an element of $\Omega^r(M)$ is defined by,

$$*(dx^{\mu_1} \wedge \cdots \wedge dx^{\mu_r}) = \frac{\sqrt{|g|}}{r!(m-r)!} g^{\mu_1\nu_1} \cdots g^{\mu_r\nu_r} \varepsilon_{\nu_1 \cdots \nu_r \nu_{r+1} \cdots \nu_m} dx^{\nu_{r+1}} \wedge \cdots \wedge dx^{\nu_m}, \quad (125)$$

in which $g^{\mu\nu}$ is the metric of manifold M , g is the determinant of $g_{\mu\nu}$, and $\mu_i, \nu_i = 1, \dots, m$.

We consider two independent cases; an r -form in $m+1$ dimensions which is an r -form along m -dimensional manifold, and an r -form in $m+1$ dimensions which is an $(r-1)$ -form along m -dimensional manifold and an 1-form along the $(m+1)$ -th direction. For the first case $\eta_r = \frac{1}{r!} \eta_{\mu_1 \cdots \mu_r} dx^{\mu_1} \wedge \cdots \wedge dx^{\mu_r}$, thus,

$$*_m \eta_r = \frac{\sqrt{|g|}}{r!(m-r)!} \eta_{\mu_1 \cdots \mu_r} g^{\mu_1\nu_1} \cdots g^{\mu_r\nu_r} \varepsilon_{\nu_1 \cdots \nu_r \nu_{r+1} \cdots \nu_m} dx^{\nu_{r+1}} \wedge \cdots \wedge dx^{\nu_m}, \quad (126)$$

$$*_{m+1} \eta_r = \frac{\sqrt{|g'|}}{r!(m+1-r)!} \eta_{\mu_1 \cdots \mu_r} g^{\mu_1\nu_1} \cdots g^{\mu_r\nu_r} \varepsilon_{\nu_1 \cdots \nu_r \nu_{r+1} \cdots \nu_m \nu_{m+1}} dx^{\nu_{r+1}} \wedge \cdots \wedge dx^{\nu_m} \wedge dx^{\nu_{m+1}}, \quad (127)$$

where g and g' are determinant of metrics in m and $m+1$ dimensional manifolds respectively, $\varepsilon_{1 \cdots m}$ is the totally antisymmetric tensor, $\mu_i = 1, \dots, m$, $\nu_i = 1, \dots, m$ for the first equation, and $\nu_i = 1, \dots, m$ for the second equation. Since the $(m+1)$ -th direction is perpendicular to the other directions one obtains $g^{\mu_i, m+1} = \delta_{\mu_i, m+1}$ and $g = g'$. Therefore in Eq. (127) ν_i only takes value in the range $1 \leq \nu_i \leq m$ for $i = 1, \dots, r+1$. Since there are $(m+1-r)$ positions for ν_{m+1} in of $\varepsilon_{1 \cdots m}$, one obtains,

$$*_{m+1} \eta_r = \frac{\sqrt{|g|}}{r!(m-r)!} \eta_{\mu_1 \cdots \mu_r} g^{\mu_1\nu_1} \cdots g^{\mu_r\nu_r} \varepsilon_{\nu_1 \cdots \nu_r \nu_{r+1} \cdots \nu_m m+1} dx^{\nu_{r+1}} \wedge \cdots \wedge dx^{\nu_m} \wedge dx^{m+1}, \quad (128)$$

where $\mu_i, \nu_i = 1 \cdots m$. By comparing Eq. (126) with Eq. (128) one obtains the following relation,

$$*_{m+1} \eta_r = *_m \eta_r \wedge dx^{m+1}. \quad (129)$$

In the second case, we assume,

$$\eta_{r+1} = \frac{1}{(r+1)!} \eta_{\mu_1 \cdots \mu_r m+1} dx^{\mu_1} \wedge \cdots \wedge dx^{\mu_r} \wedge dx^{m+1} = \frac{1}{r+1} \eta'_{rm+1} \wedge dx^{m+1}, \quad (130)$$

where $\eta'_{rm+1} \equiv \frac{1}{r!} \eta_{\mu_1 \cdots \mu_r m+1} dx^{\mu_1} \wedge \cdots \wedge dx^{\mu_r}$ is an $(r+1)$ -form along the $(m+1)$ -dimensional manifold and an r -form along the m -dimensional manifold. Hodge star operation in the m -manifold gives,

$$\begin{aligned} *_m \eta_{r+1} &= \frac{1}{r+1} *_m (\eta'_{rm+1}) \wedge dx^{m+1} \\ &= \frac{1}{r+1} \frac{\sqrt{|g|}}{r!(m-r)!} \eta_{\mu_1 \cdots \mu_r m+1} g^{\mu_1\nu_1} \cdots g^{\mu_r\nu_r} \varepsilon_{\nu_1 \cdots \nu_r \nu_{r+1} \cdots \nu_m} dx^{\nu_{r+1}} \wedge \cdots \wedge dx^{\nu_m} \wedge dx^{m+1}, \end{aligned} \quad (131)$$

where $\mu_i, \nu_i = 1 \cdots m$. In addition,

$$\begin{aligned}
*_ {m+1} \eta_{r+1} &= \frac{\sqrt{|g'|}}{(r+1)!(m-r)!} \eta_{\mu_1 \cdots \mu_r m+1} \\
&\quad g^{\mu_1 \nu_1} \cdots g^{\mu_r \nu_r} g^{m+1 \nu} \varepsilon_{\nu_1 \cdots \nu_r \nu_{r+2} \cdots \nu_{m+1}} dx^{\nu_{r+2}} \wedge \cdots \wedge dx^{\nu_{m+1}} \\
&= \frac{\sqrt{|g|}}{(r+1)!(m-r)!} \eta_{\mu_1 \cdots \mu_r m+1} \\
&\quad g^{\mu_1 \nu_1} \cdots g^{\mu_r \nu_r} g^{m+1, m+1} \varepsilon_{\nu_1 \cdots \nu_r, m+1, \nu_{r+2} \cdots \nu_{m+1}} dx^{\nu_{r+2}} \wedge \cdots \wedge dx^{\nu_{m+1}} \\
&= (-)^{m-r} \frac{\sqrt{|g|}}{(r+1)!(m-r)!} \eta_{\mu_1 \cdots \mu_r m+1} \\
&\quad g^{\mu_1 \nu_1} \cdots g^{\mu_r \nu_r} \varepsilon_{\nu_1 \cdots \nu_r \nu_{r+2} \cdots \nu_{m+1} m+1} dx^{\nu_{r+2}} \wedge \cdots \wedge dx^{\nu_{m+1}}. \quad (132)
\end{aligned}$$

Comparing Eq. (131) with Eq. (132) one verifies that

$$(*_{m+1} \eta_{r+1}) \wedge dx^{m+1} = (-)^{(m-r)} *_m \eta_{r+1}. \quad (133)$$

If the signature of $(m+1)$ -th direction is Lorentzian, then $g^{\mu_i, m+1} = -\delta_{\mu_i, m+1}$ and one should consider an extra (-1) in the above equation.

As an application of these relations one can show that for an r -form η_r along 6 dimensions of original 7-dimensional manifold,

$$\begin{aligned}
d_7^\dagger d_7 \eta_r &= (-)^{7(r+2)+1} *_7 d_7 *_7 d_7 \eta_r \\
&= (-)^{r+1} *_7 d_7 *_7 (d_6 \eta_r + dt \wedge \dot{\eta}_r) \\
&= (-)^{r+1} *_7 d_7 (*_6 d_6 \eta_r \wedge dt + *_6 \dot{\eta}_r) \\
&= (-)^{r+1} *_7 (d_6 *_6 d_6 \eta_r \wedge dt + d_6 *_6 \dot{\eta}_r + dt \wedge *_6 \ddot{\eta}_r) \\
&= (-)^{r+1} ((-)^r *_6 d_6 *_6 d_6 \eta_r + *_6 d_6 *_6 \dot{\eta}_r \wedge dt + *_6 *_6 \ddot{\eta}_r) \\
&= d_6^\dagger d_6 \eta_r - \ddot{\eta}_r + (-)^r d_6^\dagger \dot{\eta}_r \wedge dt, \quad (134)
\end{aligned}$$

where Eq. (3,4) are used to this derivation. Also for $\eta_{r+1} = \eta_r \wedge dt$, one obtains,

$$\begin{aligned}
d_7^\dagger d_7 (\eta_r \wedge dt) &= (-)^{7(r+2)+1} *_7 d_7 *_7 d_7 (\eta_r \wedge dt) \\
&= (-)^{r+1} *_7 d_7 *_7 (d_6 \eta_r \wedge dt) \\
&= *_7 d_7 (*_6 d_6 \eta_r) \\
&= *_7 (d_6 *_6 d_6 \eta_r + dt \wedge *_6 d_6 \dot{\eta}_r) \\
&= *_6 d_6 *_6 d_6 \eta_r \wedge dt + *_6 *_6 d_6 \dot{\eta}_r \\
&= -d_6^\dagger d_6 \eta_r \wedge dt + (-)^{r+1} d_6 \dot{\eta}_r. \quad (135)
\end{aligned}$$

References

- [1] E. Witten, *Chern-Simons Gauge Theory as a String Theory*, Prog.Math. 133 (1995) 637 [hep-th/9207094].
- [2] M. Bershadsky, V. Sadov, *Theory of Kähler Gravity*, Int.J.Mod.Phys. A11 (1996) 4689-4730 [hep-th/9410011].
- [3] M. Bershadsky, S. Cecotti, H. Ooguri and C. Vafa, *Kodaira-Spencer Theory of Gravity and Exact Results for Quantum String Amplitudes*, Commun. Math. Phys. 165 (1994) 311 [hep-th/9309140].
- [4] M. Bershadsky, S. Cecotti, H. Ooguri and C. Vafa, *Holomorphic Anomalies in Topological Field Theories*, Nucl.Phys. B405 (1993) 279 [hep-th/9302103].
- [5] E. Witten, *Quantum background independence in string theory*, hep-th/9306122.
- [6] E. Verlinde, *Attractors and the Holomorphic Anomaly*, hep-th/0412139.
- [7] A. A. Gerasimov and S. L. Shatashvili, *Towards Integrability of Topological Strings I: Three-forms on Calabi-Yau manifolds*, JHEP 0411 (2004) 074 [hep-th/0409238].
- [8] S. Cecotti, and C. Vafa, *Topological antitopological fusion*, Nucl. Phys. B367(1991)359.
- [9] F. Loran, *Kähler Quantization of $H^3(CY^3, R)$ and the Holomorphic Anomaly*, JHEP 0512 (2005) 004 [hep-th/0510163].
- [10] C. Vafa, *Superstrings and topological strings at large N* , J. Math. Phys. 42 (2001) 2798 [hep-th/0008142].
- [11] P. Candelas and X. Ossa, *Moduli space of Calabi-Yau Manifolds*, Nucl.Phys. B355 (1991) 455.
- [12] K. Hori, S. Katz, A. Klemm, R. Pandharipande, R. P. Thomas, C. Vafa, R. Vakil, and E. Zaslow, *Mirror symmetry*, vol. 1 of Clay Mathematics Monographs. American Mathematical Society, Providence, RI, 2003, ISBN 0821829556.
- [13] N. Hitchin, *Stable forms and special metrics*, math.DG/0107101.
- [14] R. Dijkgraaf, S. Gukov, A. Neitzke, and C. Vafa, *Topological M-theory as Unification of Form Theories of Gravity*, Adv.Theor.Math.Phys.9:593-602,2005 [hep-th/0411073].
- [15] N. Nekrasov, H. Ooguri and C. Vafa, *S-duality and topological strings*, JHEP 0410:009,2004 [hep-th/0403167].
- [16] A. Neitzke and C. Vafa, *$N=2$ strings and the twistorial Calabi-Yau*, [hep-th/0402128].
- [17] E. M. Prodanov and S. Sen, *Abelian Duality*, Phys.Rev. D62 (2000) 045009 [hep-th/9906143].
- [18] M. Nakahara, “Geometry, Topology and Physics”, IOP Publishing Ltd, 1990, ISBN 0852740948.

- [19] E. Witten, “Abelian Duality” in “Quantum Fields and Strings : A Course for Mathematicians” vol 2, Edited by P. Deligne et al, AMS, 1999, ISBN-13: 978-0-8218-2014-8.
- [20] P. Griffiths and J. Harris, “Principles of Algebraic Geometry”, Wiley, 1978, ISBN 0471327921.
- [21] A. Strominger, *Special Geometry*, Comm.Math.Phys.133 (1990) 163.
- [22] B. Hatfield, “Quantum Field Theory of Point Particles and Strings”, Addison-Wesley, 1992, ISBN 0201360799.